

Title	On Certain Nor-Continuous Functions and Shape (Shapeと無限次元多様体)
Author(s)	FUKAISHI, HIROO
Citation	数理解析研究所講究録 (1979), 342: 33-38
Issue Date	1979-01
URL	http://hdl.handle.net/2433/104289
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

On certain non-continuous functions and shape

Hiroo Fukaishi

In the shape category of topological spaces a shape morphism is constructed by a system of maps (= continuous functions) ; it is, in general, not generated by a single map.

Hence we have the following questions :

Question 1. Is it possible that a kind of non-continuous function induces a shape morphism ?

Question 2. Can a shape equivalence be generated by a certain non-continuous function ?

Definition 1. Let X and Y be topological spaces. A function $f : X \rightarrow Y$ is a connectivity function if for any connected $C \subset X$, the graph $G(f|C)$ of $f|C$ is connected.

Definition 2. A function $f : X \rightarrow Y$ is almost continuous if for any open set $N \subset X \times Y$ containing $G(f)$ there is a continuous function $g : X \rightarrow Y$ such that $G(g) \subset N$.

These notions have been considered to generalize Brouwer's fixed point theorem (cf. Stallings[10]).

Each of the following is intermediate to answer the questions.

Proposition 1. Let $f : X \rightarrow Y$ be an almost continuous function between compact metric spaces. Then there are ANR-sequences $\underline{X} = \{X_i, p_{ij}\}$ and $\underline{Y} = \{Y_i, q_{ij}\}$ with limits X and Y , respectively, and a system $\underline{f} : \underline{X} \rightarrow \underline{Y}$ of almost continuous functions $f_i : X_i \rightarrow Y_i$ such that $f_i p_{ij} = q_{ij} f_j$ for $i \leq j$.

Proof. There are ANR-sequences \underline{X} and \underline{Y} with limits X and Y , respectively, and each projection p_i, q_i surjective, since both X and Y are compact metric. Define a function $f_i : X_i \rightarrow Y_i$ for each i , by the formula $f_i p_i = q_i f$. The almost continuity of f implies that of f_i .

Proposition 2. A bijective connectivity function with connectivity inverse function does not induce a shape equivalence.

Proof. By the example of Stallings [10,p.262]. Let X be the circle represented as the real numbers mod 1. Define a function $f : X \rightarrow X \times X$ by the formula

$$f(x \bmod 1) = 1/x \bmod 1, \text{ where } 0 < x \leq 1.$$

Let Y be the graph of f and $f^* : X \rightarrow Y$ such that

$$f^*(x) = (x, f(x)).$$

Then f^* is a bijection, and both f^* and f^{*-1} are connectivity functions, but $\text{Sh}(X) \neq \text{Sh}(Y)$, because their 1-dimensional Čech cohomology groups are different.

REFERENCES ON NON-CONTINUOUS FUNCTIONS

- [1] Noguchi, H., A generalization of absolute neighborhood retracts, *Kôdai Math. Sem. Rep.*, 5(1953), 20-22.
- [2] Fan, K. & Struble, R.A., Continuity in terms of connectedness, *Indag. Math.*, 16(1954), 161-164.
- [3] Tanaka, T., On the family of connected subsets and the topology of spaces, *J. of Math. Soc. Japan*, 7(1955), 389-393.
- [4] Nash, J., Generalized Brouwer theorem, *Bull. Amer. Math. Soc.*, 62(1956), 76, (Research problem).
- [5] Hamilton, O.H., Fixed points for certain noncontinuous transformations, *Proc. Amer. Math. Soc.*, 8(1957), 750-756.
- [6] O'Neill, B. & Straus, E.G., A fixed point theorem, *Proc. Amer. Math. Soc.*, 8(1957), 1148-1151.
- [7] O'Neill, B., Induced homology homomorphisms for set-valued maps, *Pacific J. Math.*, 7(1957), 1179-1184.
- [8] Pervin, W.J. & Levine, N., Connected mappings of Hausdorff spaces, *Proc. Amer. Math. Soc.*, 9(1958), 488-496.
- [9] Ponomarev, V.I., A new space of closed sets and many-valued mappings, *D. A. N. SSSR*, 118(1958), 1081-1084.
- [10] Stallings, J., Fixed point theorems for connectivity maps, *Fund. Math.*, 47(1959), 249-263.
- [11] Long, P.E., Properties of certain non-continuous transformations, *Duke Math. J.*, 28(1961), 639-645.
- [12] Hildebrand, S.K. & Sanderson, D.E., Connectivity functions and retracts, *Fund. Math.*, 57(1965), 237-245.

- [13] Roberts, J.H., Zero-dimensional sets blocking connectivity functions, *Fund. Math.*, 57(1965), 173-179.
- [14] Cornette, J.L., Connectivity functions and images on Peano continua, *Fund. Math.*, 58(1966), 183-192.
- [15] Hagan, M.R., Equivalence of connectivity maps and peripherally continuous transformations, *Proc. Amer. Math. Soc.*, 17(1966), 175-177.
- [16] Naimpally, S.A., Graph topology for function spaces, *Trans. Amer. Math. Soc.*, 123(1966), 267-272.
- [17] Whyburn, G.T., Connectivity of peripherally continuous functions, *Proc. Nat. Acad. Sci. U. S. A.*, 55(1966), 1040-1041.
- [18] Cornette, J.L. & Girolo, J.E., Connectivity retracts of finitely coherent Peano continua, *Fund. Math.*, 61(1967), 177-182.
- [19] Poppe, H., Über Graphentopologien für Abbildungsräume, I, *Bull. Acad. Polon. Sci.*, 40(1967), 71-80.
- [20] Fuller, R.V., Relations among continuous and various non-continuous functions, *Pacific J. Math.*, 25(1968), 495-509.
- [21] Gmurczyk, A., On approximative retracts, *Bull. Acad. Polon. Sci.*, 16(1968), 9-14.
- [22] Granas, A., Fixed point theorems for the approximative ANR-s, *Bull. Acad. Polon. Sci.*, 16(1968), 15-19.
- [23] Hyman, D.M., ANR divisors and absolute neighborhood contractibility, *Fund. Math.*, 62(1968), 61-73.
- [24] Lin, S.T., Fixed point properties and inverse limit spaces, *Pacific J. Math.*, 25(1968), 117-122.

- [25] Hyman, D.M., On decreasing sequences of compact absolute retracts, *Fund. Math.*, 64(1969), 91-117.
- [26] Hoyle, III, H.B., Connectivity maps and almost continuous functions, *Duke Math. J.*, 37(1970), 671-680.
- [27] Naimpally, S.A. & Pareek, C.M., Graph topologies for function spaces, II, *Annales Soc. Math. Polon. Ser. I.*, 13(1970), 221-231.
- [28] Brown, J.B., Totally discontinuous connectivity functions, *Colloq. Math.*, 23(1971), 53-60.
- [29] Clapp, M.H., On a generalization of absolute neighborhood retracts, *Fund. Math.*, 70(1971), 117-130.
- [30] Gmurczyk, A., Approximative retracts and fundamental retracts, *Colloq. Math.*, 23(1971), 61-63.
- [31] Chewning, W.C., Connectivity retracts of unicoherent Peano continua in \mathbb{R}^n , *Fund. Math.*, 75(1972), 25-27.
- [32] Kellum, K.R. & Garrett, B.D., Almost continuous real functions, *Proc. Amer. Math. Soc.*, 33(1972), 181-184.
- [33] Powers, M.J., Fixed point theorems for non-compact approximative ANR's, *Fund. Math.*, 75(1972), 62-68.
- [34] Kellum, K.R., Almost continuous functions on I^n , *Fund. Math.*, 79(1973), 213-215.
- [35] Bogaty, S.A., Approximative and fundamental retracts, *Math. USSR Sbornik*, 22(1974), 91-103.
- [36] Felt, J.E., ϵ -continuity and shape, *Proc. Amer. Math. Soc.*, 46(1974), 426-430.
- [37] Kellum, K.R., Sums and limits of almost continuous functions, *Colloq. Math.*, 31(1974), 125-128.

- [38] Cannon, J. W. , Taming cell-like embedding relations, Lecture Notes in Math., 439, Springer, (1975), 66-118.
- [39] Kellum, K. R. , On a question of Borsuk concerning non-continuous retracts I, Fund. Math., 87(1975), 90-92.
- [40] Kellum, K. R. , Non-continuous retracts, Stavrakas & Allen ed., Studies in topology, Academic Press, (1975), 255-261.
- [41] Miller, Jr., M. H. , Discontinuous G_δ graphs, *ibid.*, 383-392.
- [42] Kellum, K. R. , On a question of Borsuk concerning non-continuous retracts II, Fund. Math., 92(1976), 135-140.
- [43] Kellum, K. R. , The equivalence of absolute almost continuous retracts and ε -absolute retracts, Fund. Math., 96(1977), 229-235.
- [44] Ranchin, D. V. , On modification of topologies in connection with the given discontinuous mappings of the space into itself, Bull. Acad. Polon. Sci., 26(1978), 149-156.